

# FIESTA®

## GRANULAR WEED & FEED

PRE- AND POST-EMERGENT 8-0-1 HERBICIDE

6 in 1!

Fertilizes your turf while controlling weeds, moss, disease\*, algae\* and lichens!

\*not for use in California

Active Ingredient	By weight
Sodium Ferric EDTA	14.0%
Other Ingredients	86.0%
Total	100.0%

EPA Registration No. 67702-63

EPA Establishment No. 9198-OH-1M, 9198-OH-2B, 9198-AL-1A, 9198-IN-3S, 9198-NC-1N

First digit of batch code indicates establishment number.

NET WEIGHT 20 LBS

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency information concerning this product, call the poison control center at 1-800-222-1222.

**Environmental Hazards:** For terrestrial uses. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all applicable directions and precautions on this label before using.

**SITES:** Lawns, turf (use on rights of way or non-crop areas), golf courses, parks, playgrounds, cemeteries, athletic fields.

### Pests:

Controls or suppresses weeds such as but not limited to:

- Black medic (*Medicago lupulina*)§
- Broadleaf plantain (*Plantago major*)\*
- Bull thistle (*Cirsium vulgare*)
- Canada thistle (*Cirsium arvense*)\*
- Common chickweed (*Stellaria media*)§
- Crabgrass (*Digitaria spp.*)- pre-emergent control only
- Creeping buttercup (*Ranunculus repens*)
- Creeping Charlie/Ground ivy (*Glechoma hederacea*)\*
- Dandelion (*Taraxicum officinale*)
- Dovefoot geranium (*Geranium molle*)\*
- English daisy (*Bellis perennis*)§
- False dandelion (*Hypochoeris radicata*)\*
- Heal-all (*Prunella vulgaris*)\*
- Lawn burweed (*Soliva ptero-sperma*)\*
- Narrowleaf plantain (*Plantago lanceolata*)\*
- Oxalis (*Oxalis spp.*)\*
- Prostrate knotweed (*Polygonum aviculare*)\*

- Shepherd's-purse (*Capsella bursa-pastoris*)\*
- Silverweed cinquefoil (*Potentilla anserina*)\*
- Speedwell spp. (including Persian Speedwell (*Veronica persica*) and Slender speedwell (*Veronica filiformis*)\*)
- White clover (*Trifolium repens*)
- Wild chamomile (*Matricaria chamomilla*)\*
- Wild geranium (*Geranium robertianum*)\*
- moss, algae\*, lichens\*, liverworts\*

Controls or suppresses diseases such as but not limited to:

- Dollar Spot\*
- Rust\*
- Snow Mold\*

\* Not for use in California

§ only approved for suppression in California

**Application Directions:** Apply at a rate of 3-8 lbs/1000 ft<sup>2</sup> (0.5 - 1.15 oz/yard<sup>2</sup>), or 5-8 lbs/1000 ft<sup>2</sup> (0.72 - 1.15 oz/yard<sup>2</sup>) in California. The standard rate for this product is 5 lbs/1000 ft<sup>2</sup> (0.72 oz/yard<sup>2</sup>); use the higher rate of 8 lbs/1000 ft<sup>2</sup> (1.15 oz/yard<sup>2</sup>) on tough perennial weeds, such as common chickweed§, buttercup and broadleaf plantain\*. Use at 3 lbs/1000 ft<sup>2</sup>† (0.5 oz/yard<sup>2</sup>) on young moss, and weed seedlings of false dandelion\*, English daisy\*, dandelion\* or white clover\*. Repeat treatment in 3 to 4 weeks for best results.

For Spot Treatment, apply 5 lbs/1000 ft<sup>2</sup>. Repeat treatment in 3 to 4 weeks for best results.

For Pre-emergent control of broadleaf weeds and crabgrass, begin application when soil temperatures reach 50 to 55°F. Apply 5-8 lbs/1000 ft<sup>2</sup>, depending on weed pressure. Repeat treatment in 3 to 6 weeks for best results.

Application rate	Weeds	Diseases	Area Treated
3 lbs/1000 ft <sup>2</sup> (0.43 oz/yd <sup>2</sup> )†	Weed seedlings of English daisy* ( <i>Bellis perennis</i> ), dandelion* ( <i>Taraxacum officinale</i> ), false dandelion* ( <i>Hypochoeris radicata</i> ), young moss.	Dollar spot*, snow mold*, and rust*. Use the higher rate when disease pressure is high.	6670 ft <sup>2</sup> /20 lb container
4 lbs/1000 ft <sup>2</sup> (0.58 oz/yd <sup>2</sup> )†	Moss, speedwell species, including Persian speedwell ( <i>Veronica persica</i> ), and slender speedwell* ( <i>Veronica filiformis</i> ), wild geranium* ( <i>Geranium robertianum</i> ).		5000 ft <sup>2</sup> /20 lb container
5 lbs/1000 ft <sup>2</sup> (0.72 oz/yd <sup>2</sup> )	Black medic ( <i>Medicago lupulina</i> ) §, bull thistle ( <i>Cirsium vulgare</i> ), Canada thistle* ( <i>Cirsium arvense</i> ), creeping charlie/ground ivy* ( <i>Glechoma hederacea</i> ), dandelion ( <i>Taraxacum officinale</i> ), dove's foot geranium* ( <i>Geranium molle</i> ), English daisy ( <i>Bellis perennis</i> ), false dandelion* ( <i>Hypochoeris radicata</i> ), heal-all* ( <i>Prunella vulgaris</i> ), lawn burweed* ( <i>Soliva ptero-sperma</i> ), narrowleaf plantain* ( <i>Plantago lanceolata</i> ), oxalis* ( <i>Oxalis spp.</i> ), prostrate knotweed* ( <i>Polygonum aviculare</i> ), Shepherd's-purse* ( <i>Capsella bursa-pastoris</i> ), silverweed cinquefoil* ( <i>Potentilla anserina</i> ), algae*, lichens*, liverworts*. Pre-emergent control of Crabgrass ( <i>Digitaria spp.</i> ).		4000 ft <sup>2</sup> /20 lb container
8 lbs/1000 ft <sup>2</sup> (1.15 oz/yd <sup>2</sup> )	Common chickweed§ ( <i>Stellaria media</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), broadleaf plantain* ( <i>Plantago major</i> ), white clover* ( <i>Trifolium repens</i> ).		2500 ft <sup>2</sup> /20 lb container

† Lower application rate not for use in California. Minimum application rate in California is 5 lbs/1000 ft<sup>2</sup>

\* Not for use in California

§ only approved for suppression in California

### How to Apply:

Contents may have settled during shipment; invert bag several times to ensure product is mixed.

For best results, apply a few days after mowing when temperature is between 50°F and 85°F. If temperature is below 80°F ensure turf is wet from dew, or water turf lightly before application. If temperature is above 80°F do not pre-wet grass and apply product in the late afternoon or evening. Do not exceed the 5 lbs/1000 ft<sup>2</sup> rate when temperatures are above 85°F. Do not apply when daytime temperature exceeds 90°F.

Apply using a broadcast/rotary, drop, or handheld spreader set to the correct application rate. Uniform coverage is important to prevent turf damage. See table below for spreader settings. Wait at least 24 hours after application to water turf. Turf areas can be re-sown one day after treatment.

Apply product only on turf. Application on hard surfaces can cause staining. If product gets on surface, sweep off if the surface is dry, rinse off immediately if the surface is wet. If product gets on clothing or footwear, wash immediately.

### Spreader settings:

Weed Species	Rate	Scotts® Broadcast and Rotary Spreader Setting (All Models)	Scotts® Drop Spreader Setting (All Models)	Scotts® Hand-Held Spreader Setting (All models)
Weed seedlings of English daisy* ( <i>Bellis perennis</i> ), dandelion* ( <i>Taraxacum officinale</i> ) false dandelion* ( <i>Hypochoeris radicata</i> ), young moss.	3 lbs per 1000 square feet	2.5	5	3
Black medic ( <i>Medicago lupulina</i> )§, bull thistle ( <i>Cirsium vulgare</i> ), Canada thistle* ( <i>Cirsium arvense</i> ), creeping charlie/ground ivy* ( <i>Glechoma hederacea</i> ), dandelion ( <i>Taraxacum officinale</i> ), dove's foot geranium* ( <i>Geranium molle</i> ), English daisy ( <i>Bellis perennis</i> ), false dandelion* ( <i>Hypochoeris radicata</i> ), heal-all* ( <i>Prunella vulgaris</i> ), lawn burweed* ( <i>Soliva ptero-sperma</i> ), narrowleaf plantain* ( <i>Plantago lanceolata</i> ), oxalis* ( <i>Oxalis spp.</i> ), prostrate knotweed* ( <i>Polygonum aviculare</i> ), Shepherd's-purse* ( <i>Capsella bursa-pastoris</i> ), silverweed cinquefoil* ( <i>Potentilla anserina</i> ), speedwell species, including Persian speedwell, ( <i>Veronica persica</i> ) and slender speedwell* ( <i>Veronica filiformis</i> ), wild geranium* ( <i>Geranium robertianum</i> ), moss, algae*, lichens*, liverworts*. Preemergent control of crabgrass ( <i>Digitaria spp.</i> ).	5 lbs/1000 square feet	3	6	3.5
Common chickweed§ ( <i>Stellaria media</i> ), creeping buttercup ( <i>Ranunculus repens</i> ), broadleaf plantain* ( <i>Plantago major</i> ), white clover* ( <i>Trifolium repens</i> ).	8 lbs/1000 square feet	4.5	7.5	5

† Lower application rate not for use in California. Minimum application rate in California is 5 lbs/1000 ft<sup>2</sup>

\* Not for use in California

§ only approved for suppression in California

Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions. Spreaders other than those listed in this table may be used, as long as they are calibrated prior to use to deliver the product within the use rates on this label.

### What will I see?

Works quickly; dying weeds will be seen within 24 hours of application. Treated weeds, moss, or algae\* will turn brown or black. Occasionally a darkening of the grass leaf blades can occur after treatment; however, the grass will recover within a few days to weeks. May stain thatch (dead grass under turf); this does not affect the health of the grass. If turf is damaged, irrigation 24 hours after application, followed by mowing to remove the turf tip burn can help turf recover faster.

\* Not for use in California

### Use Precautions:

Do not apply to drought stressed grass; ensure turf is well watered prior to application. Do not apply when the daytime temperature will exceed 90°F. Do not apply to bentgrass. If unsure of grass sensitivity, make a test application to a small area of the turf. Avoid applying to desirable plants. Do not apply if rainfall is expected within 24 hours of application. Do not irrigate within 24 hours of application.

### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Keep this product in its original container and in a secured area that is inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Plastic containers: Triple rinse or equivalent. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Paper and plastic bags: Completely empty into application equipment. Then dispose of bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BATCH CODE:

### 8-0-1

#### Guaranteed Analysis

Total nitrogen (N) 8.0% Other Water Soluble Nitrogen	8.0%	Derived from soy protein hydrolysate, potassium chloride, iron EDTA.
Soluble Potash (K <sub>2</sub> O)	1.0%	
Iron (Fe) 1.8% Chelated Iron (Fe)	1.8%	Chlorine (Cl), maximum 1.0%

**In PA, NJ, NH, MD, VA:** Do not exceed 0.7 lb. of readily available nitrogen or 0.9 lb. of total nitrogen per 1,000 sq. ft. per application. Do not exceed 3.2 total nitrogen per 1,000 sq. ft. per year.

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfc.org/metals.html>.

FL License No. F003462

Florida Applications: We recommend that you follow the Florida Green Industries BMP's at: <http://www.flrules.org/Gateway/reference.asp?No=Ref-04706>

We recommend that you follow the Florida Golf Course BMP's at: <http://www.flrules.org/Gateway/reference.asp?No=Ref-04702>

• **Fiesta Granular Weed & Feed is formulated to feed your turf and control broadleaf weeds, moss, algae\*, diseases\* commonly found in turf.**

• **Pre- and Post-emergent activity of broadleaf weeds with one application.**

• **Builds thick, green turf while killing listed weeds.**

• **Treats up to 6670 ft<sup>2</sup>**

• **Works fast.**

• **Weeds start dying within 24 hours.**

• **Fast and effective.**

• **Spring and fall weed control.**

• **Rainfast in 24 hours.**

• **No unpleasant odor.**

• **Ready-to-use dry formulation.**

Patent Pending.

### NOTICE TO BUYER

Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. To the extent consistent with applicable law, this warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller. To the extent consistent with applicable law, the Seller shall not be liable for any such use. To the extent consistent with applicable law, the Seller makes no other warranties, either expressed or implied.

### Registrant:

**W. Neudorff GmbH KG**  
An der Muhle 3,  
31860 Emmerthal, Germany  
[www.neudorffpro.com](http://www.neudorffpro.com)

